

Methods and Tips to Use Aqueous (Water-Based) IBA Rooting Solutions[®]

Joel Kroin

Hortus USA Corp., P.O. Box 1956, New York, New York 10113, USA

Email: support@hortus.com

Since the discovery of natural plant rooting hormones for propagation (Thimann and Went, 1934), it has been intuitive to apply these substances to the basal end of cuttings to produce new roots. Basal methods used are the Quick Dip and Basal Long Soak Methods, using rooting solutions, and the Dry Dip Method, using rooting hormone powders.

Currently, many growers use foliar applied aqueous IBA rooting solutions to propagate annual, perennial and woody plants. Dr. Fred Davies' physiology research proved efficacy of the application (Davies, 1978, 1980, 1982). Since indole-3-butyric acid (IBA) is a natural rooting hormone (Epstein, 1984, 1993), having such ability is understandable. When aqueous IBA solutions are applied to the leaves of cuttings, the IBA enters the plant's vascular system through open stomata. Indole-3-butyric acid then translocates, by polar transport, to the basal end; there, plants store it for self-regulated root formation.

Kees Eigenraam (1985) developed commercial foliar methods, first used by Dutch chrysanthemum growers. Kees and I standardized the techniques, the foliar Spray Drip Down and Total Immerse Methods. In the early 1990s we introduced the methods to USA growers, first for annual and perennial plants. Research by Bailey Nurseries expanded uses to woody plants (Drahn, 2007). Required for foliar methods, aqueous IBA rooting solutions are made using Rhizopon[®] AA Water Soluble Tablets and Hortus IBA Water Soluble Salts[®].

METHODS

The following is a brief review of the basal and foliar methods along with some "tips." Some of the studies and tips are in the Combined Proceedings of the IPPS (Kroin 1992, 2008, 2009, 2010, 2011) and *Hortus Plant Propagation from Cuttings* (Kroin, 2011a).

Basal Methods

1. Basal Quick Dip Method. Hydrate cuttings. Using rooting solutions, dip the basal end of the cuttings about $\frac{3}{4}$ in. into the solution for a few seconds. Stick in propagation medium. Take care of cuttings.

2. Basal Long Soak Method. Hydrate cuttings. Batch treat, using rooting solutions, soak the basal end of the cuttings, about 1 in. into the solution for several hours (sometimes 12-48 h). Stick in propagation medium. Take care of cuttings.

3. Basal Dry Dip Method. Hydrate cuttings. Using dry rooting hormone powders, dip the basal end of the cuttings about $\frac{3}{4}$ in. into the dry powder. Stick in propagation medium. Take care of cuttings.

Foliar Methods

1. Spray Drip Down Method. Stick cuttings into propagation medium. Hydrate cuttings. Batch treat, using aqueous IBA rooting solutions, spray onto the leaves until visually the droplets fall. Turn on misters after about $\frac{3}{4}$ h or until the solution dries on the leaves. Take care of cuttings.

2. Total Immerse Method. Hydrate cuttings. Batch treat, using aqueous IBA rooting solutions, total immerse the cuttings a few seconds, drain. Stick cuttings into propagation medium. Take care of cuttings.

TIPS

The studies listed used solutions made with Hortus IBA Water Soluble Salts[®]. Alternatively it is possible to use Rhizopon AA Water Soluble Tablets (Distributor: Phytotronics. <Sales@phytotronics.com>, phytotronics.com. 314-330-0717; Rhizopon

worldwide: Rhizopon bv, rhizopon.com).

Foliar Aqueous IBA Rooting Solution Treatment

1. Timing for the Spray Drip Down Method. Using aqueous IBA rooting solutions spray the leaves of the cuttings soon after sticking. It is best to treat cuttings on the same day as sticking. Waiting to spray several days after sticking gives reduced root formation. (Kroin, 2010; Davies, 1978, 1980, 1982). If after sticking the cuttings are under stress from high daytime air temperatures in the propagation area, keep the cuttings hydrated. Some growers spray early the following morning after sticking, when air temperatures are in the range from 65-90°F.

2. Aqueous Versus Alcohol-Based IBA Solutions. Studies found using the Spray Drip Down Method that alcohol in the IBA rooting solution caused cutting death (Kroin, 2011). This is likely due to alcohol solutions dehydrating cells. Alcohol-based solutions if used by Quick Dip can cause what growers call “alcohol burns” with subsequent cutting failure. For foliar methods only use aqueous IBA rooting solutions.

3. Spray Drip Down Method Treatment in Warm versus Cold Temperature. Studies compared the Spray Drip Down Method using aqueous IBA rooting solutions with solution and cutting temperatures at 45 and 78°F. For cuttings treated and maintained at 45°F for 2 h, then brought to 78°F, had deaths. For cuttings treated and maintained at 78°F had improved root formation as compared with 78°F control cuttings (Kroin, 2011).

It is best to do foliar methods when the cutting and solution temperatures are nominally in the range from 65-90°F.

4. For Hard-to-Root Cuttings and Unseasonal Rooting, Comparison of the Basal Long Soak Method with the Total Immerse Method. The Basal Long Soak Method successfully roots cuttings of many kinds (Thimann, 1950). Rooting studies were performed on *Buxus sinensis* cuttings in May. It is not the usual time to propagate these cuttings. Cuttings treated by Basal Long Soak Method had more roots than those treated by the Total Immerse Method. Basal Long Soak treated cuttings had more fine roots than any other cuttings. Totally immersed cuttings had more roots than untreated control cuttings (Kroin, 2011). Since the Basal Long Soak Method allows slow absorption in the plant's vascular system, it may be an effective method when other ways are not successful.

5. Wetting Agents Are not Needed in Foliar Applied Aqueous IBA Rooting Solutions. Spray Drip Down Method studies found wetting agents added to the aqueous IBA rooting solution had no noticeable difference in rooting compared with solutions at the same IBA concentration that had no added wetting agent (Kroin, 2011).

6. Foliar Rates Are Low. When doing foliar method trials, use low rates of aqueous IBA rooting solutions. Typical trial rates for foliar methods:

- Annual plants: 50-250 ppm IBA.
- Perennials and woody plants: 250, 500, 1000, 1500 ppm IBA, where 1000 ppm is often the selected rate.
- Rates above 1500 ppm are rarely needed.

7. Treatment Area. Using the Spray Drip Down Method, 1 gal can treat an area of 175-225 ft² of plants (for growth stimulation) or cuttings (for propagation).

8. Mistors. When using the Spray Drip Down Method, after sticking, keep the cuttings well hydrated. Turn off misters, spray treat, and wait about 45 min or until the spray solution dries then restore misting.

9. Aqueous IBA Rooting Solutions. Aqueous IBA rooting solutions can and are used by many applications where IBA and K-IBA are specified, usually without regard to prior use of alcohol based solutions.

10. Conversion from Technical IBA to Hortus IBA Water Soluble Salts or Rhizopon AA Water Soluble Tablets. Many articles relating to plant propagation and growing specify use of IBA or K-IBA rooting solutions. In most cases, aqueous IBA can be used. To make aqueous IBA rooting solutions, growers use Hortus IBA Water Soluble Salts or Rhizopon AA Water Soluble Tablets. They are U.S.A. EPA registered for this use. In

rooting solution, they are as – if made with K-IBA. Note: Under the federal law FIFRA, the USA. EPA prohibits use of unregistered end-use technical grade IBA and K-IBA for plant growing. When converting rates from “technical” IBA or K-IBA, my advice to growers is to trial Hortus IBA Water Soluble Salts or the Tablets about 10% lower rate than technical. “Technical” products may not be labeled as true assay, or the salts and tablets have enhanced activity due to formulation.

11. Make Stock Solutions. Making concentrated Hortus IBA Water Soluble Salts aqueous IBA stock solutions avoids the need to measure the salt powder often. Solutions can be made to over 100,000 ppm IBA. Make a solution in multiples of the number of grams per unit volume; mark the side of the container into that number of multiples. When making a working solution, decant the stock solution to the appropriate mark and pour it into the working tank. Next, add water to the working to bring to the required volume.

12. Quality of Water and Temperature to Make Solutions. Use clean water to make aqueous IBA rooting solutions with Hortus IBA Water Soluble Salts or Rhizopon AA Water Soluble Tablets. Hard water may cause some precipitation. Demineralized water is best. Do not use pond water; it likely contains biological components that will breakdown the IBA. Hortus IBA Water Soluble Salts or Rhizopon AA Water Soluble Tablets easily dissolve in water at 65-95°F. They dissolve most easily at the higher range of temperatures.

13. Dispose of Solutions Dipped with Plant Material. The foliar Total Immerse, Basal Quick Dip, and Basal Long Soak Methods require plants to be dipped into the working solution. Avoid pathogen cross contamination by disposing of the solution between lots.

14. Secondary Aqueous IBA Rooting Solution Applications. The Spray Drip Down Method is useful for secondary applications to cuttings that are already stuck and had been untreated or treated by any method, either basal or foliar. Uses are to level crops, induce new root formation and improve root mass. Spray the aqueous IBA rooting solution on the cuttings at the same rate as the original useful rate or lower.

15. Typical Trial Rates for Secondary Applications

- Annual plants: 50-200 ppm IBA.
- Perennials and woody plants: 250-1,500 ppm IBA.

Treating Tissue Culture Plantlets

Tissue culture plantlets have poorly functioning stomata or few in number. The Total Immerse Method has proven successful for transplant treatment. Before planting, totally immerse the plantlets in the aqueous rooting solution for approximately 5 sec. Rates used are Rhizopon AA Water Soluble Tablets at 1-3 tablets per liter of water.

CONCLUSIONS

Several basal and foliar auxin application methods can be used to propagate plants from cuttings. Based upon trials, growers can find one or more methods that are suitable for their needs based upon the plant variety and species, seasonal variations, production needs, and total cost. When rooting solutions are selected, aqueous IBA rooting solutions are useful by all methods.

Literature Cited

- Davies, Jr., F.T. 1978. Histological and physiological analysis of adventitious root formation *Ficus primula*. A Dissertation Presented to the Graduate Council of The University of Florida.
- Davies, Jr., F.T. 1980. Growth regulator effects on adventitious root formation in leaf bud cuttings of juvenile and mature *Ficus pumila*. Amer. Soc. Hort. Sci. 105(1):91-95.
- Davies, Jr., F.T. 1982. Initiation and development of roots in juvenile and mature leaf bud cuttings of *Ficus pumila* L. Amer. J. Bot. 69(5):804-811.
- Davies, Jr., F.T. 1982. Shoot RNA, cambial activity, and indolebutyric acid effectivity in seasonal rooting of juvenile and mature *Ficus pumila* cuttings. Physiol. Plant. 62:571-

575.

- Davies, Jr., F.T. 1988. The physiological basis of adventitious root formation. *Acta Hort.* 227:113-120.
- Drahn, S. 2007. Auxin application via foliar sprays. *Comb. Proc. Intl. Plant Prop. Soc.* 57:274-277.
- Eigenraam, K. 2011. Current recommendations for use of Rhizopon rooting hormones. *Comb. Proc. Intl. Plant Prop. Soc.* 61:187-191.
- Epstein, E. and Lavee, S. 1984. Conversion of indole-3-butyric acid to indole-3-acetic acid by cuttings of grapevine (*Vitis vinifera*) and olive (*Olea europea*). *Plant Cell Physiol.* 25:697-703.
- Epstein, E. and Ludwig-Muller, J. 1993. Indole-3-butyric acid in plants: occurrence, synthesis, metabolism and transport. *Physiol. Plant.* 88:382-389.
- Hartman, H., Kester, D., Davies, Jr., F.T. and Geneve, R. 2010. *Plant Propagation Principles and Practices*. 8th edition. Prentice Hall, Upper Saddle River, New Jersey 07458.
- Kroin, J. 1992. Advances using Indole-3-butyric acid (IBA) dissolved in water for rooting cuttings, transplanting and grafting. *Comb. Proc. Intl. Plant Prop. Soc.* 42:489-492.
- Kroin, J. 2008. Propagate plants from cuttings using dry-dip rooting powders and water based rooting solutions. *Comb. Proc. Intl. Plant Prop. Soc.* 58:360-372.
- Kroin, J. 2009. Propagation of plants from cuttings using rooting solutions by foliar methods. *Comb. Proc. Intl. Plant Prop. Soc.* 59:437-453.
- Kroin, J. 2010. Propagation of cuttings using foliar applied IBA in aqueous solutions at or after sticking. *Comb. Proc. Intl. Plant Prop. Soc.* 60:369-377.
- Kroin, J. 2011. How to improve cutting propagation using water based indole-3-butyric acid rooting solutions. *Comb. Proc. Intl. Plant Prop. Soc.* 61:381-391.
- Kroin, J. 2011a. Hortus plant propagation from cuttings. A guide to using plant rooting hormones. Hortus U.S.A., PO Box 1956, New York, New York 10113. support@hortus.com
- Thimann, K. and Went, F. 1934. On the chemical nature of root forming hormone. *Proc. Royal Acad. Amsterdam.* 38:456-459.
- Thimann, K. and Behnke-Rogers, J. 1950. *The Use of Auxins in the Rooting of Woody Cuttings*. Harvard Forest, Persham, Massachusetts.